

## Real Analysis Qual, Spring 2020

**Problem 1.** Prove that if  $f : [0, 1] \rightarrow \mathbb{R}$  is continuous then

$$\lim_{k \rightarrow \infty} \int_0^1 kx^{k-1} f(x) \, dx = f(1).$$

*Proof.* First, for  $0 \leq x < 1$ , we claim  $kx^{k-1} \rightarrow 0$  pointwise. If  $x = 0$ , then the claim is obvious. Otherwise,  $x^{-k+1}$  tends to infinity in  $k$ . Therefore, by L'Hopital's Rule, we have

$$\lim_{k \rightarrow \infty} kx^{k-1} = \lim_{k \rightarrow \infty} \frac{k}{x^{-k+1}} = \lim_{k \rightarrow \infty} \frac{1}{(-k+1)x^{-k}} = 0.$$

Moreover,  $kx^{k-1}$  is monotone. So, on the interval  $[0, a]$  for  $a < 1$ ,  $kx^{k-1}$  is bounded by  $ka^{k-1}$ . Therefore,  $kx^{k-1} \rightarrow 0$  uniformly on  $[0, a]$ .

Now,  $f$  is a continuous function on the compact interval  $[0, 1]$ , and so  $f$  is bounded by some constant  $M$ . Then, for all  $a \in [0, 1)$ ,

$$\int_0^a kx^{k-1}(-M) \, dx \leq \int_0^a kx^{k-1} f(x) \, dx \leq \int_0^a kx^{k-1} M \, dx.$$

Since  $kx^{k-1} \rightarrow 0$  uniformly on  $[0, a]$ , we obtain  $\lim_{k \rightarrow \infty} \int_0^a kx^{k-1} f(x) \, dx = 0$  for each  $a$ . So, for all  $a \in [0, 1)$ , we have

$$\lim_{k \rightarrow \infty} \int_0^1 kx^{k-1} f(x) \, dx = \lim_{k \rightarrow \infty} \int_a^1 kx^{k-1} f(x) \, dx.$$

Define  $N_a$  to be the lower bound of  $f(x)$  and  $M_a$  to be the upper bound of  $f(x)$  over the compact interval  $[a, 1]$ . Then,

$$N_a - a^k N_a = \int_a^1 kx^{k-1} N_a \, dx \leq \int_a^1 kx^{k-1} f(x) \, dx \leq \int_a^1 kx^{k-1} M_a \, dx = M_a - a^k M_a.$$

Since  $a^k \rightarrow 0$ , then taking  $k \rightarrow \infty$ , we have for all  $a \in [0, 1)$ ,

$$N_a \leq \lim_{k \rightarrow \infty} \int_a^1 kx^{k-1} f(x) \, dx = \lim_{k \rightarrow \infty} \int_0^1 kx^{k-1} f(x) \, dx \leq M_a.$$

Finally, by continuity of  $f$ , as  $a \rightarrow 1$ , we have  $N_a \rightarrow f(1)$  and  $M_a \rightarrow f(1)$ . Therefore,

$$f(1) \leq \lim_{k \rightarrow \infty} \int_0^1 kx^{k-1} f(x) \, dx \leq f(1).$$

So, we obtain equality, as needed. □

**Problem 2.** Let  $m_*$  denote the Lebesgue outer measure on  $\mathbb{R}$ .

- (a) Prove that for every  $E \subseteq \mathbb{R}$  there exists a Borel set  $B$  containing  $E$  with the property that

$$m_*(B) = m_*(E).$$

- (b) Prove that if  $E \subseteq \mathbb{R}$  has the property that  $m_*(A) = m_*(A \cap E) + m_*(A \cap E^c)$  for every set  $A \subseteq \mathbb{R}$ , then there exists a Borel set  $B \subseteq \mathbb{R}$  such that  $E = B \setminus N$  with  $m_*(N) = 0$ .

*Proof.* We begin with (a). By definition, if  $E \subseteq \mathbb{R}$ , then

$$m_*(E) = \inf \left\{ \sum_{n=1}^{\infty} V(I_n) : (I_n)_{n=1}^{\infty} \text{ are open-closed intervals, } E \subseteq \bigcup_{n=1}^{\infty} I_n \right\},$$

where  $V((a, b]) = b - a = m_*(I_n)$ . From definition of infimum, there exists a sequence  $(I_n^{(k)})_{n=1}^{\infty}$  such that  $E \subseteq \bigcup_{n=1}^{\infty} I_n^{(k)}$  and  $\sum_{n=1}^{\infty} m_*(I_n^{(k)}) \leq m_*(E) + 1/k$ . Define  $B_k = \bigcup_{n=1}^{\infty} I_n^{(k)}$ . Then,  $B_k$  is a countable union of open-closed intervals, and thus Borel. Set  $B = \bigcap_{k=1}^{\infty} B_k$  and note that  $B$  is Borel. For each  $k$ , we have  $B_k \supseteq E$ , and so  $B \supseteq E$ . Moreover, for all  $k$ , we have  $m_*(E) \leq m_*(B) \leq m_*(B_k) \leq m_*(E) + 1/k$ . Thus,  $m_*(E) = m_*(B)$  for  $B$  Borel.

For (b), take  $E$  as defined. By (a), we choose  $B$  such that  $B$  is Borel,  $E \subseteq B$ , and  $m_*(E) = m_*(B)$ . Then, since  $E \setminus B = \emptyset$ , we have

$$0 = m_*(E \setminus B) = m_*((B \setminus E) \cap E) + m_*((B \setminus E) \cap E^c) = m_*(\emptyset) + m_*(B \cap E^c) = m_*(B \cap E^c).$$

Moreover,  $B = (B \cap E) \cup (B \cap E^c) = E \cup (B \cap E^c)$ . Taking  $N = B \cap E^c$ , we have  $E = B \setminus N$  with  $m_*(N) = 0$  for  $B$  a Borel set.  $\square$

### Problem 3.

- (a) Prove that if  $f \in L^1(\mathbb{R})$ , then

$$\lim_{N \rightarrow \infty} \int_{|x| \geq N} |f(x)| \, dx = 0$$

and demonstrate that it is not necessarily the case that  $f(x) \rightarrow 0$  as  $N \rightarrow \infty$ .

- (b) Prove that if  $f \in L^1([1, \infty))$  and decreasing, then  $\lim_{x \rightarrow \infty} f(x) = 0$  and in fact  $\lim_{x \rightarrow \infty} x f(x) = 0$ .
- (c) If  $f : [1, \infty) \rightarrow [0, \infty)$  is decreasing with  $\lim_{x \rightarrow \infty} x f(x) = 0$ , does this ensure  $f \in L^1([1, \infty))$ ?

*Proof.* We start with (a). First, suppose  $f$  is a compactly supported continuous function. Then,  $\text{supp } f$  is bounded, say by  $M$ . Therefore, for all  $|x| \geq M$ , we have  $f(x) = 0$ . So, for all  $N \geq M$ , we obtain

$$\int_{|x| \geq N} |f(x)| \, dx \geq \int_{|x| \geq M} |f(x)| \, dx = \int_{|x| \geq M} 0 \, dx = 0.$$

Therefore,

$$\lim_{N \rightarrow \infty} \int_{|x| \geq N} |f(x)| \, dx = 0.$$

Now, say that  $f \in L^1(\mathbb{R})$ . Continuous functions with compact support are dense in  $L^1(\mathbb{R})$ . Let  $\epsilon > 0$ . Choose  $g$  to be a continuous function with compact support such that  $\|f - g\|_1 < \epsilon$ . Then, take  $N$  large enough so that  $\int_{|x| \geq N} |g| dx = 0$ . We have

$$\int_{|x| \geq N} |f(x)| dx \leq \int_{|x| \geq N} |f(x) - g(x)| dx + \int_{|x| \geq N} |g(x)| dx \leq \int_{\mathbb{R}} |f(x) - g(x)| dx < \epsilon.$$

So,

$$\lim_{N \rightarrow \infty} \int_{|x| \geq N} |f(x)| dx = 0.$$

Now, define  $f_N = \sum_{n=1}^N \mathbb{1}_{[2^n, 2^{n+1}/2^n]}$ . Then,  $f_N \rightarrow f_\infty$  pointwise. Moreover,  $f_N \leq f_{N+1}$ , so the  $f_N$  are monotone. Finally, the  $f_N$  are all nonnegative. Thus, by MCT, we have

$$1 = \sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{N \rightarrow \infty} \sum_{n=1}^N \frac{1}{2^n} = \lim_{N \rightarrow \infty} \sum_{n=1}^N \int \mathbb{1}_{[2^n, 2^{n+1}/2^n]} dx = \lim_{N \rightarrow \infty} \int f_N dx = \int f_\infty dx.$$

Since  $|f_\infty| = f_\infty$ , then  $f_\infty \in L^1(\mathbb{R})$ . But,  $\lim_{x \rightarrow \infty} f_\infty(x) \neq 0$ , since for all  $n$  we have  $f_\infty(2^n) = 1$ .

Now we prove (b). Say that  $f(x)$  is monotone in  $x$ . Say there is some  $x_0 \in [1, \infty)$  such that  $f(x_0) < 0$ . Then, for all  $x \geq x_0$ , we have  $f(x) \leq f(x_0)$ . Therefore,

$$\int_{x_0}^{\infty} f(x) dx \geq \int_{x_0}^{\infty} f(x_0) dx = -\infty.$$

So,  $\int_1^{\infty} |f| dx = \infty$ , a contradiction. Thus,  $f(x) \geq 0$  for all  $x$ . Therefore,  $f(x)$  decreases monotonically in  $x$  and is bounded below, so  $\lim_{x \rightarrow \infty} f(x) = L$  for some  $L$ . Say that  $L > 0$ . Then, there exist constants  $M, N$  such that for all  $x \geq N$  we have  $f(x) \geq M > 0$ . Then,

$$\int_N^{\infty} |f(x)| dx = \int_N^{\infty} f(x) dx \geq \int_N^{\infty} M dx = \infty,$$

again a contradiction. Thus,  $L = 0$ . Furthermore, fix  $x$ . Then,

$$\int_{x/2}^x f(t) dt \geq \int_{x/2}^x f(x) dt = f(x) \Big|_{x/2}^x = x f(x) - \frac{x}{2} f(x) = \frac{x f(x)}{2}.$$

Now, since the tail of the integral of an  $L^1$  function goes to 0, then  $\lim_{x \rightarrow \infty} \int_{x/2}^x f(t) dt = 0$ .

Therefore,  $\lim_{x \rightarrow \infty} \frac{x f(x)}{2} = 0$ , so  $\lim_{x \rightarrow \infty} x f(x) = 0$ .

Finally, we prove (c). Consider

$$f(x) = \frac{1}{(x+1) \ln(x+1)}$$

which is defined on all  $[1, \infty)$ . Then,  $x f(x) = \frac{x}{(x+1) \ln(x+1)} \rightarrow_{x \rightarrow \infty} 0$ . Moreover,  $f(x)$  is decreasing, since  $(x+1) \ln(x+1)$  increases monotonically on  $[1, \infty)$ . Finally, the antiderivative of  $f(x)$  on  $[1, \infty)$  is  $\ln(\ln(x+1))$ . Since  $\ln(\ln(x+1)) \rightarrow \infty$  as  $x \rightarrow \infty$ , and  $\ln(\ln(2))$  is a constant, then

$$\infty = \int_1^{\infty} f(x) dx = \int_1^{\infty} |f(x)| dx.$$

So  $f(x) \notin L^1([1, \infty))$ . □

**Problem 4.** Let  $f \in L^1(\mathbb{R})$  and  $g \in L^1(\mathbb{R})$ . Argue that  $H(x, y) = f(y)g(x - y)$  defines a function in  $L^1(\mathbb{R}^2)$  and deduce from this that

$$f * g(x) = \int f(y)g(x - y) \, dy$$

defines a function in  $L^1(\mathbb{R})$  that satisfies

$$\|f * g\|_1 \leq \|f\|_1 \|g\|_1.$$

*Proof.* Since  $f, g$  are measurable functions, and  $(x, y) \mapsto xy$  is a continuous function, then  $H(x, y)$  is a measurable function. By Tonelli's Theorem, we obtain

$$\begin{aligned} \int |H(x, y)| \, dm_2 &= \int \int |f(y)g(x - y)| \, dx \, dy \\ &= \int |f(y)| \int |g(x - y)| \, dx \, dy \\ &= \int |f(y)| \int |g(x)| \, dx \, dy \\ &= \left( \int |f(y)| \, dy \right) \left( \int |g(x)| \, dx \right). \end{aligned}$$

Therefore,  $H(x, y) \in L^1(\mathbb{R}^2)$ . Furthermore,

$$\int |f * g(x)| \, dx = \int \left| \int f(y)g(x - y) \, dy \right| \, dx \leq \int \int |f(y)g(x - y)| \, dy \, dx = \|H\|_1.$$

So,  $f * g \in L^1(\mathbb{R})$ , and

$$\|f * g\|_1 \leq \|H\|_1 = \int |H(x, y)| \, dm_2 = \left( \int |f(y)| \, dy \right) \left( \int |g(x)| \, dx \right) = \|f\|_1 \|g\|_1,$$

completing the proof. □

**Problem 5.** Compute the following limit and justify your calculations:

$$\lim_{n \rightarrow \infty} \int_0^n \left( 1 + \frac{x^2}{n} \right)^{-n+1} \, dx.$$

*Proof.* Recall that  $\lim_{n \rightarrow \infty} (1 + \frac{y}{n})^n \rightarrow e^y$  pointwise. Likewise,  $1 + \frac{x^2}{n} \rightarrow 1$  pointwise. Thus, for  $x$  fixed we have

$$\lim_{n \rightarrow \infty} \mathbb{1}_{[0, n]} \left( 1 + \frac{x^2}{n} \right)^{-n+1} = \lim_{n \rightarrow \infty} \frac{1 + \frac{x^2}{n}}{\left( 1 + \frac{x^2}{n} \right)^n} = \lim_{n \rightarrow \infty} \frac{1}{\left( 1 + \frac{x^2}{n} \right)^n} = e^{-x^2}.$$

Moreover, for each  $n$ , applying the Binomial Theorem and observing that  $x \geq 0$ , we have the inequality

$$\left( 1 + \frac{x^2}{n} \right)^n = \sum_{k=0}^n \binom{n}{k} \left( \frac{x^2}{n} \right)^k \geq \binom{n}{2} \left( \frac{x^2}{n} \right)^2 = \frac{n-1}{2n} x^4 \geq \frac{1}{4} x^4.$$

Therefore,

$$\left(1 + \frac{x^2}{n}\right)^{-n+1} = \frac{1 + \frac{x^2}{n}}{\left(1 + \frac{x^2}{n}\right)^n} \leq \frac{4\left(1 + \frac{x^2}{n}\right)}{x^4} \leq \frac{4 + 4x^2}{x^4}.$$

Furthermore, on the interval  $[0, 1]$ , since  $1 + x^2/n \geq 1$  for all  $x \in [0, 1]$ , we have

$$\left(1 + \frac{x^2}{n}\right)^{-n+1} = \frac{1}{\left(1 + \frac{x^2}{n}\right)^{n-1}} \leq 1.$$

Define

$$f(x) := \begin{cases} 1, & \text{if } x \in [0, 1], \\ \frac{4+4x^2}{x^4}, & \text{if } x \in (1, \infty). \end{cases}$$

By our previous arguments,  $f(x)$  globally bounds  $\left(1 + \frac{x^2}{n}\right)^{-n+1}$ . Moreover,

$$\int_0^\infty |f(x)| dx = \int_0^1 f(x) dx + \int_1^\infty f(x) dx = \int_0^1 1 dx + \int_1^\infty \frac{4 + 4x^2}{x^4} dx < \infty.$$

So,  $\left(1 + \frac{x^2}{n}\right)^{-n+1}$  has the integrable dominant  $f$ . Therefore, by DCT, we obtain

$$\lim_{n \rightarrow \infty} \int_0^n \left(1 + \frac{x^2}{n}\right)^{-n+1} dx = \int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2},$$

given that  $\int_0^\infty e^{-x^2} dx$  is the standard Gaussian integral. □

### Problem 6.

- (a) Show that  $L^2([0, 1]) \subseteq L^1([0, 1])$  and that  $\ell^1(\mathbb{Z}) \subseteq \ell^2(\mathbb{Z})$ .
- (b) For  $f \in L^1([0, 1])$  define

$$\hat{f}(n) := \int_0^1 f(x) e^{-2\pi i n x} dx.$$

Prove that if  $f \in L^1([0, 1])$  and  $\{\hat{f}(n)\} \subseteq \ell^1(\mathbb{Z})$ , then

$$S_N f(x) = \sum_{|n| \leq N} \hat{f}(n) e^{2\pi i n x}$$

converges uniformly on  $[0, 1]$  to a continuous function  $g$  that equals  $f$  almost everywhere.

*Hint: One possible approach is to argue that if  $f \in L^1([0, 1])$  with  $\{\hat{f}(n)\} \in \ell^1(\mathbb{Z})$ , then  $f \in L^2([0, 1])$ .*

*Proof.* We begin with (a). Let  $f$  be an  $L^2([0, 1])$  function. Then, by Hölder's Inequality, we have

$$\|f\|_1 \leq \|f\|_2 \|1\|_2 = \|f\|_2 < \infty.$$

So,  $f \in L^1([0, 1])$ . Now, say that  $(a_n) \in \ell^1(\mathbb{Z})$ . Since  $\mathbb{Z}, \mathbb{N}$  are both countable, we may simply assume this series is indexed in  $\mathbb{N}$ . Then,  $a_n \rightarrow 0$ . So, pick  $N$  such that for all  $n \geq N$ , we have  $|a_n| < 1$ . Then,  $|a_n|^2 \leq |a_n|$ , so we obtain

$$\sum_{n=1}^{\infty} |a_n|^2 = \sum_{n=1}^N |a_n|^2 + \sum_{n=N}^{\infty} |a_n|^2 \leq \sum_{n=1}^N |a_n|^2 + \sum_{n=1}^{\infty} |a_n| < \infty.$$

Therefore,  $(a_n) \in \ell^2(\mathbb{Z})$ .

We now prove (b). Take  $M \geq N$ . Then,

$$\|S_N f - S_M f\|_1 = \left\| \sum_{N \leq |n| \leq M} \widehat{f}(n) e^{-2\pi i n x} \right\| \leq \sum_{N \leq |n| \leq M} \int \left| \widehat{f}(n) e^{-2\pi i n x} \right| dx = \sum_{N \leq |n| \leq M} |\widehat{f}(n)|.$$

Since  $(\widehat{f}(n)) \in \ell^1(\mathbb{Z})$ , then we conclude that  $S_N f$  is Cauchy. Since  $L^1([0, 1])$  is complete, then  $S_N f$  converges to some  $g \in L^1([0, 1])$ . Moreover, observe that  $|\widehat{f}(n) e^{-2\pi i n x}| \leq |\widehat{f}(n)|$  for all  $x$ . Then, since  $\sum_{n \in \mathbb{Z}} |\widehat{f}(n)| < \infty$ , by the Weierstrass M-test, we have a uniform convergence  $S_N f \rightarrow g$ . Since each  $S_N f$  is continuous, then  $g$  is continuous. Now, since  $S_N f \rightarrow g$  uniformly, then we may pass the limit through the integral. That is,

$$\widehat{g}(n) = \int_0^1 g(x) e^{-2\pi i n x} dx = \lim \int_0^1 S_N f(x) e^{-2\pi i n x} dx = \lim \int_0^1 \widehat{f}(n) dx = \widehat{f}(n),$$

by the orthogonality of  $(e^{2\pi i n x})_{n \in \mathbb{Z}}$  over  $L^2([0, 1])$ . Therefore,  $f$  and  $g$  have the same Fourier coefficients. By uniqueness of fourier coefficients, we conclude that  $f = g$  a.e..  $\square$